

**DRUG FACTS**

Active ingredient(s)

Purpose

**Use(s)**  
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

**Warnings**

**Do not use**  
if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**  
liver or kidney disease. Your doctor should determine if you need a different dose.

**Ask a doctor or pharmacist before use if you are**

**When using this product**  
do not take more than directed. Taking more than directed may cause drowsiness.

**Stop use and ask a doctor if**  
an allergic reaction to this product occurs. Seek medical help right away.

**If pregnant or breast-feeding**

**Keep out of reach of children**  
In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

adults and children 6 years and over	1 tablet daily; not more than 1 tablet in 24 hours
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

**Other information**

- do not use if blister unit is broken or torn (Blister Only)
- do not use if printed foil under cap is broken or missing (Bottle Only)
- store at 20°-25°C (68°-77°F)
- protect from excessive moisture (Blister Only)

**Inactive ingredients**  
lactose monohydrate, magnesium stearate, povidone, pregelatinized starch